arXiv:1403.7960v1 [hep-ph] 31 Mar 2014

Search for HZZ' couplings at the LHC

V. Arı* and O. Çakır†

Ankara University, Department of Physics, 06100, Ankara, Turkey

S. Kuday‡

Ankara University, Institute of Accelerator Technologies, 06700, Ankara, Turkey

Abstract

New physics models predict the possibility of extra neutral gauge bosons (Z') associated with an extra $U(1)'$ gauge symmetry. We study the couplings of the Higgs boson to the Z boson and Z' boson predicted by the new physics models. The couplings of the Z' boson to quarks can also be investigated through the $Z'q\bar{q}$ interactions. The accessible ranges of the parameter space have been searched for processes $pp \to HZX$ and $pp \to HHZX$ at the LHC with $\sqrt{s} = 14$ TeV.

*Electronic address: vari@science.ankara.edu.tr

†Electronic address: ocakir@science.ankara.edu.tr

‡Electronic address: kuday@science.ankara.edu.tr

I. INTRODUCTION

The experimental opportunities can be exploited to search for hints beyond the standard model (BSM) physics after the recent discovery of the Higgs boson with a mass of 125 GeV by the ATLAS and CMS experiments [1, 2] at the LHC. Due to its gauge charges the Higgs boson may interact with the BSM fields. This new interaction can also modify the couplings between the Higgs and SM fields at tree level or loop level. Within some extensions of the standard model (SM) a new neutral gauge boson Z' can be included [3–5].

The phenomenological studies on the Z' boson signatures can be found in [6–8]. The present constraints on the Z' mass and couplings from both electron-positron and hadron-hadron colliders have been presented in [9]. The precision measurements at the Z boson resonance lead to the limit on $Z - Z'$ mixing [10]. The ATLAS and CMS experiments at the LHC, running at higher center of mass energy and luminosity, have updated the Tevatron limits on the Z' boson mass [11, 12]. The ATLAS Collaboration searches for a massive resonance decaying to top quark pairs in the hadronic channels, and excludes the leptophobic Z' boson with mass smaller than 1.32 TeV [14]. The CMS Collaboration excludes leptophobic Z' boson resonance with a mass of $m_{Z'} < 1.3$ TeV for its width $\Gamma_{Z'} = 0.012 m_{Z'}$ in the search of heavy resonance decaying into $t\bar{t}$ pair with subsequent leptonic decays [13].

In this work, we investigate the HZZ' couplings via the processes $pp \to HZX$ and $pp \to HHZX$ at the LHC with $\sqrt{s} = 14$ TeV. The couplings of the Z' boson to quarks can also be investigated via $Z'q\bar{q}$ interactions. For the signal process $pp \to HZX$, the Z' boson contributes in the s-channel resonance diagrams through family diagonal neutral current couplings. However, the signal process $pp \to HHZX$ has contributions from both the Z'-boson resonance and the Z-exchange diagrams. We obtain the accessible ranges of the mass for two different Z' models at the four-fermion and six-fermion final states resulting from $H \to b\bar{b}$ and $Z \to l^+l^-$ decays. These modes exist even if the Z' boson decouples from the leptons.

II. THE PRODUCTION CROSS SECTION

The interaction of the Higgs boson (H) with the Z boson and the new Z' boson can be written through the kinetic term of the scalar field. The relevant interaction term can be

Table I: Vector and axial-vector couplings of Z' boson predicted by leptophobic (LP) model and η (ETA) model.

	C_V^u	C_A^u	C_V^d	C_A^d	C_V^l	C_A^l	C_V^ν	C_A^ν
Z'(LP)	$-7/9$	1	$2/9$	0	0	0	0	0
Z'(ETA)	0	$4/6$	$1/2$	$1/6$	$-1/2$	$1/6$	$-1/6$	$-1/6$

written as $L_{HZZ'} = -g_Z g_{Z'} z[H] \upsilon H Z_\mu Z'^\mu$. The new boson Z' couples to the fermion field (f) with the interaction term $L_{f\bar{f}Z'} = (g_{Z'}/2)\bar{f}\gamma^\mu(C_V^f - C_A^f\gamma^5)f$. Here, the vector ($C_V^f$) and axial-vector ($C_A^f$) couplings are model dependent. We use a linear change of variable $g_{Z'} = \gamma_{Z'} g_e/\cos\theta_W \sin\theta_W$ for the coupling constant. The triple coupling HZZ' depends on the $U(1)'$ charge. In the model the triple coupling parameter $z[H] = -1$. The $U(1)'$ couplings to the fermions for two different Z' models are given in Table I. Since the extensively studied Drell-Yan process even constrained the possible $Z'l^+l^-$ couplings, we consider two candidate models in which the Z' bosons have small or no couplings to leptons.

The production processes have contribution from the resonance production of Z' boson. Therefore, we already included the decay width of the Z' boson in the calculation within different Z' models, as shown in Fig. 1 for leptophobic (LP) and η (ETA) models. The branching ratios versus Z' mass are given in Fig. 2 and Fig. 3 for the LP and ETA models, respectively.

At the LHC with $\sqrt{s} = 14$ TeV, the cross section of the HZ production through the signal process is about 0.20 pb and 0.15 pb with Z' boson mass $m_{Z'} = 1.5$ TeV for the LP and ETA model, respectively. The cross section of HHZ production is around 3.85 fb and 2.78 fb with Z' boson mass $m_{Z'} = 1.5$ TeV for the LP and ETA model, respectively. As shown in Figs. 4 - 5 and 6 - 7, the background cross sections result in 0.7 pb and 0.29 pb for HZ and HHZ production at $\sqrt{s} = 14$ TeV, respectively. The final states of $2l + 2b_{jet}$ and $2l + 4b_{jet}$ are the main backgrounds for the HZ and HHZ productions. In order to suppress the relevant background, we can apply the invariant mass cuts, on dileptons to be around the Z boson mass, on dijets with each jets as b-tagged to be around the Higgs mass.



Figure 1: Decay width of Z' boson depending on its mass for two different models.



Figure 2: Branching ratios of the Z' boson as predicted by lepto-phobic model (LP) depending on its mass.



Figure 3: Branching ratios of the Z' boson as predicted by η (ETA) model of E_6 depending on its mass.



Figure 4: The cross section for the process $pp \to HZX$ depending on the mass of Z' boson for LP model.



Figure 5: The cross section for the process $pp \to HZX$ depending on the mass of Z' boson for ETA model.



Figure 6: The cross section for the process $pp \to HHZX$ depending on the mass of Z' boson for LP model.



Figure 7: The cross section for the process $pp \to HHZX$ depending on the mass of Z' boson for ETA model.

III. PARAMETER SPACE ANALYSIS

In the analysis, we use the cross section of the signal and background for the final states explained in the previous section. A pair of b-jets give an invariant mass distributions peak around the Higgs mass and the dilepton invariant mass distribution give peak around the Z mass. It is also helpful to use angular seperations of the leptons, b-jets and between leptons and jets. The massive Z' boson can be reconstructed from the resonance peak in the invariant mass spectrum of the final states $2l + 2b_{jet}$ and $2l + 4b_{jet}$ originating from the HZ and HHZ productions, respectively. The luminosity needs for 3σ signal observability for final state $2l + 2b_{jet}$ depending on the mass of Z' boson for two different Z' models ETA and LP are given in Fig. 8. Having an integrated luminosity of $L_{int} = 100$ fb^{-1} at the LHC with $\sqrt{s} = 14$ TeV, it is possible to search for Z' boson up to a mass value of $M_{Z'} = 1.9$ TeV and 2.0 TeV for the ETA and LP models as seen from Fig. 8. The search range for the Z'boson mass can be found as $M_{Z'} = 1.5$ TeV and 1.8 TeV for ETA and LP models for $2l + 4b_{jet}$ as given in Fig. 9.



Figure 8: Luminosity requirement for 3σ signal observability for final state $2l + 2b_{jet}$ depending on the mass of Z' boson for two different Z' models ETA and LP.



Figure 9: Luminosity requirement for 3σ signal observability for final state $2l + 4b_{jet}$ depending on the mass of Z' boson for two different Z' models ETA and LP.

IV. CONCLUSIONS

We present the attainable parameter space for two different $U(1)'$ models by studying the processes $pp \to HZX$ and $pp \to HHZX$ at the LHC with $\sqrt{s} = 14$ TeV. The HZ production has large cross section and it has the advantage of two b-jets and two charged leptons in the final state. The background for HHZ production has lower cross section than HZ production, however it has the advantage of four b-jets and two charged leptons in the final state. The Higgs boson mostly decays into two b-jet channel and we use the advantage of efficient identification of dileptons from Z boson. The couplings of Z' boson to the SM quarks play a role in the production, and the $U(1)'$ charge of the Higgs boson scales the HZZ' coupling. The searches for HVV couplings will also extend our perspectives for new physics at the LHC with $\sqrt{s} = 14$ TeV and $L_{int} = 100$ fb^{-1}.

[1] G. Aad *et al.* (ATLAS Collaboration), Phys. Lett. B **716**, 1 (2012).

[2] S. Chatrchyan *et al.* (CMS Collaboration), Phys. Lett. B **716**, 30 (2012).

[3] D. London and J. L. Rosner, Phys. Rev. D **34**, 1530 (1986).

[4] A. Leike, Phys. Rept. **317**, 143 (1999).

[5] P. Langacker, Rev. Mod. Phys. **81**, 1199 (2009).

[6] V. Barger *et al.*, Phys. Lett. B **580**, 186 (2004).

[7] M.S. Carena *et al.*, Phys. Rev. D **70**, 093009 (2004).

[8] K. Agashe *et al.*, Phys. Rev. D **81**, 096002 (2010).

[9] J. Beringer *et al.*, (Particle Data Group), Phys. Rev. D **86**, 010001 (2012).

[10] V.V. Andreev and A.A. Pankov, Phys. Atom. Nucl. **75**, 76 (2012).

[11] T. Aaltonen *et al.*, (CDF Collaboration), Phys. Rev. Lett. **102**, 031801 (2009).

[12] V.M. Abazov *et al.*, (D0 Collaboration), Phys. Lett. B **695**, 88 (2011).

[13] S. Chatrchyan *et al.*, (CMS Collaboration), arXiv:1211.3338 [hep-ex].

[14] G. Aad *et al.*, (ATLAS Collaboration), arXiv:1211.2202 [hep-ex].